

The Stochastic Burgers Equation

L. Bertini^{1,2}, N. Cancrini¹, G. Jona-Lasinio^{1,3}

¹ Dipartimento di Fisica, Università di Roma “La Sapienza”, P.le Aldo Moro 2, 00185 Roma, Italy. E-mail: jona@roma1.infn.it, cancrini@sci.uniroma1.it

² Dipartimento di Matematica, Università di Roma “Tor Vergata”, Via delle Ricerca Scientifica, 00133 Roma, Italy. E-mail: bertini@mat.utovrm.it

³ Centro Linceo Interdisciplinare, Via della Lungara 10, I-00165 Roma, Italy

Received: 21 September 1993

Abstract: We study Burgers Equation perturbed by a white noise in space and time. We prove the existence of solutions by showing that the Cole-Hopf transformation is meaningful also in the stochastic case. The problem is thus reduced to the analysis of a linear equation with multiplicative *half white* noise. An explicit solution of the latter is constructed through a generalized Feynman-Kac formula. Typical properties of the trajectories are then discussed. A technical result, concerning the regularizing effect of the convolution with the heat kernel, is proved for stochastic integrals.

1. Introduction

One of the first attempts to arrive at the statistical theory of turbulent fluid motion was the proposal by Burgers of his celebrated equation

$$\partial_t u_t(x) = \nu \partial_x^2 u_t(x) - u_t(x) \partial_x u_t(x), \quad (1.1)$$

where $u_t(x)$ is the velocity field and ν is the viscosity. As Burgers emphasized in the introduction of his book [3] this equation represents an extremely simplified model describing the interaction of dissipative and non-linear inertial terms in the motion of the fluid. A clear discussion on the physical problems connected with Burgers equation can be found in [10]. As shown by Cole and Hopf [5, 7], Eq. (1.1) can be explicitly solved and, in the limit of vanishing viscosity, the solution develops shock waves.

Rigorous results have been recently established in the study of some statistical properties: random initial data are considered in [1, 14, 16], while in [15] a forcing term, which is a stationary stochastic process in time and a periodic function in space, is added.

The study of Burgers equation with a forcing term is interesting in view of the phenomenological character of (1.1). Since it represents an incomplete description of a system, a forcing term can provide a good model of the neglected effects; in particular a random perturbation may help to select interesting invariant measures.

Translational invariance is preserved when (1.1) is perturbed by additive stochastic processes stationary in space and time.

In principle one can think of a wide variety of stationary random forcing terms. White noise in time and space is very often a candidate and the main motivations behind this choice are a central limit type argument and the insufficient knowledge of the neglected effects or external disturbances. A basic feature of white noise is its singularity at small scales. This may be unphysical in certain cases, but, as stressed in [2], it seems reasonable to expect that when a white noise of small amplitude is added to a deterministic equation the effects of small scales should not be overwhelming in determining the macroscopic behaviour of the system. In other words, using a terminology from quantum field theory, the equation should exhibit some *ultraviolet stability*. We also note that with this choice, due to the absence of time correlations, the full Galilean invariance of (1.1) is preserved.

In this paper we establish an existence theorem for the Cauchy problem for Burgers equation perturbed by an additive white noise in space and time. Furthermore the theorem gives an explicit expression for the solution. In order to illustrate our result let us write the equation and fix the notations

$$\partial_t u_t(x) = \nu \partial_x^2 u_t(x) - u_t(x) \partial_x u_t(x) + \varepsilon \eta_t(x), \tag{1.2}$$

where $t \in \mathbf{R}^+$, $x \in \mathbf{R}$, ε is the noise intensity and $\eta_t(x)$ is white noise in space and time, i.e.

$$\mathbf{E}(\eta_t(x)\eta_{t'}(x')) = \delta(t - t')\delta(x - x'). \tag{1.3}$$

We realize the white noise $\eta_t(x)$ as the generalized derivative of the brownian sheet, i.e. $\eta_t(x) = \partial_t \partial_x W_t(x)$. The gaussian process $W_t(x)$ has correlation function

$$\mathbf{E}^w(W_t(x)W_{t'}(x')) = t \wedge t' C(x, x'), \quad C(x, x') = \theta(xx') |x| \wedge |x'|, \tag{1.4}$$

where $a \wedge b = \min\{a, b\}$ and θ is the indicator function of the set $[0, \infty)$. We remark that $C(x, x')$ is a Lipschitz function.

We write (1.2) as an integral equation using the Green's function of its linear part

$$u_t(x) = G_t * u_0(x) - \frac{1}{2} \int_0^t ds G'_{t-s} * u_s^2(x) + \varepsilon \int_0^t G'_{t-s} * dW_s(x), \tag{1.5}$$

where $*$ is the convolution with the heat kernel

$$G_t(x) = (4\nu\pi t)^{-1/2} \exp \left\{ -\frac{x^2}{4\nu t} \right\}, \tag{1.6}$$

i.e.

$$G_t * f(x) = \int dy G_t(x - y) f(y) \tag{1.7}$$

and

$$G'_t * f(x) = \int dy \partial_x G_t(x - y) f(y) \tag{1.8}$$

finally in (1.5) $dW_s(x)$ is the stochastic integral with respect to the brownian sheet and $u_0(x)$ is the initial condition. Equation (1.5) is meaningful for a wider class of functions than (1.2), i.e. for functions which do not possess two derivatives.

We shall construct a solution of Eq. (1.5) by using the Cole-Hopf [5, 7] transformation $u_t(x) = -2\nu\partial_x \log \psi_t(x)$. Proceeding formally it reduces (1.2) to the following linear equation with multiplicative *half white* noise

$$\partial_t \psi_t(x) = \nu \partial_x^2 \psi_t(x) - \frac{\varepsilon}{2\nu} \psi_t(x) \partial_t W_t(x). \tag{1.9}$$

In [12] analogous equations, but with white noise, have been studied. Their discrete version have also been considered, e.g. [11].

Burgers equation (1.2) is invariant under translation of the space variable x . Due to the presence of the *half white* noise $\partial_t W_t(x)$ this property, as can be seen from (1.4), does not hold for (1.9). We show in Sect. 4 how the translation invariance is recovered through the Cole-Hopf transformation.

In order to study rigorously (1.9), one has to interpret it as a Stochastic PDE. Since it contains a non-trivial diffusion the stochastic differential presents the well known ambiguities. In order to obtain, via the Cole-Hopf transformation, the solution of Burgers equation (1.5), the stochastic differential in (1.9) has to be interpreted in the Stratonovich sense [8] as we show in Sect. 4. In the following we thus consider

$$d\psi_t(x) = \nu \partial_x^2 \psi_t(x) dt - \frac{\varepsilon}{2\nu} \psi_t(x) \circ dW_t(x) \tag{1.10}$$

which can be written in terms of the Ito differential as

$$d\psi_t(x) = \left(\nu \partial_x^2 \psi_t(x) + \frac{\varepsilon}{2\nu} V(x) \psi_t(x) \right) dt - \frac{\varepsilon}{2\nu} \psi_t(x) dW_t(x); \tag{1.11}$$

the extra term $V(x) = \frac{1}{2} C(x, x) = \frac{1}{2} |x|$ arises from the formal expression of the Stratonovich differential, see (1.4) above. In this paper we first construct, via a generalized Feynman-Kac formula, a process which solves the mild form of (1.11),

$$\psi_t(x) = G_t * \psi_0(x) + \frac{\varepsilon}{2\nu} \int_0^t G_{t-s} * (V \psi_s ds - \psi_s dW_s)(x), \tag{1.12}$$

where $\psi_0(x)$ is the initial condition, it is related to $u_0(x)$ by $u_0(x) = -2\nu\partial_x \log \psi_0(x)$. We then prove that the Cole-Hopf transformation can be applied to $\psi_t(x)$ and it defines a solution of (1.5).

In order to give a precise formulation of our results we have to specify in greater detail the mathematical objects appearing in the previous equations.

2. Preliminaries and Results

Brownian Sheet. We denote by $(\Omega, \mathcal{F}, P^w)$ the probability space of the brownian sheet $W_t(x)$; for its properties see [17]. We introduce the filtration $\mathcal{F}_t = \sigma\{W_s(x) : (s, x) \in [0, t] \times \mathbf{R}\}$, the minimal σ -algebra such that $W_s(x)$ is measurable for all $s \leq t, x \in \mathbf{R}$.

For Eq. (1.12) we need an integral with respect to the brownian sheet which is an Ito integral in the time variable t and a Bochner integral in the space variable x . We thus define, according to Walsh's terminology, the following *martingale measure*

$$W_t(A) = \int_A W_t(x) dx, \quad t \in \mathbf{R}^+, A \in \mathcal{B}(\mathbf{R}), \tag{2.1}$$

where $\mathcal{B}(\mathbf{R})$ is the Borel σ -algebra of \mathbf{R} . As $W_t(x)$ is dP^w -a.s. continuous the above integral is well defined. For every $A, W_t(A)$ is a \mathcal{F}_t martingale; the cross variation of the martingales $W_t(A), W_t(B)$ is

$$\langle W(A), W(B) \rangle_t = t \int_{A \times B} dx dx' C(x, x'). \tag{2.2}$$

According to Walsh ([17], Chap. II) we can thus define the Ito-Bochner integral with respect to the brownian sheet for \mathcal{F}_t progressively measurable functions $f = f(t, x, \omega)$ such that

$$\mathbf{E}^w \int_0^t ds (f, Cf) = \mathbf{E}^w \int_0^t ds \int dx dx' C(x, x') f(s, x, \omega) f(s, x', \omega) < \infty. \tag{2.3}$$

This integral will be denoted by

$$I_t(f) = \int_0^t \int dx f(s, x, \omega) dW_s(x); \tag{2.4}$$

it is a \mathcal{F}_t martingale with quadratic variation

$$\langle I(f), I(f) \rangle_t = \int_0^t ds (f, Cf). \tag{2.5}$$

In the next sections we use the Burkholder-Davis-Gundy inequality (see e.g. [13], IV 4.2) for the stochastic integral (2.4): for all $p \geq 2$ exists $c_p > 0$ such that

$$\|I_t(f)\|_p \leq c_p \left\| \int_0^t ds (f, Cf) \right\|_{p/2}^{1/2}, \tag{2.6}$$

where here and throughout the paper $\|\cdot\|_p$ is the norm in $L^p(dP^w)$. The inequality (2.6) follows from the martingale property and (2.5).

Stochastic Curvilinear Integral. We shall construct a solution of (1.12) via a Feynman-Kac formula. In this formula a Stochastic Curvilinear Integral that we now define will appear. Let $s \mapsto \varphi_s$ be a Hölder continuous function from $[0, t]$ to \mathbf{R} and $s_k = 2^{-n}kt, k = 0, \dots, 2^n$ be a partition of $[0, t]$, introduce

$$M_\varphi^n(t) = \sum_k (W_{s_{k+1}}(\varphi_{s_k}) - W_{s_k}(\varphi_{s_k})); \tag{2.7}$$

using (1.4) we have

$$\lim_{n \rightarrow \infty} \mathbf{E}^w (M_\varphi^n(t))^2 = \lim_{n \rightarrow \infty} \sum_k C(\varphi_{s_k}, \varphi_{s_k})(s_{k+1} - s_k) = \int_0^t ds |\varphi_s| \tag{2.8}$$

which has the geometrical interpretation of the area under the curve $s \mapsto |\varphi_s|$. It is not difficult to verify that $M_\varphi^n(t)$ is a Cauchy sequence in $L^2(dP^w)$; its limit defines

$$M_\varphi(t) = \int_0^t dW_s(\varphi_s), \quad t \in [0, T]. \tag{2.9}$$

$M_\varphi(t)$ is a dP^w -a.s. continuous gaussian process and a \mathcal{F}_t martingale. If $s \mapsto \gamma_s$ is another curve the cross variation of $M_\varphi(t)$ and $M_\gamma(t)$ is

$$\langle M_\varphi, M_\gamma \rangle_t = \int_0^t ds C(\varphi_s, \gamma_s). \tag{2.10}$$

We note that a weaker form of the integral (2.9) was introduced, in a different context, in [4]. A general theory of stochastic curvilinear integrals is developed in [6]; there the Lipschitz property of the process in the x variable is assumed, therefore our case is not included in that theory.

Results. On the initial condition $u_0(x)$ we assume the following. It is a continuous function we write in the form $u_0(x) = -\partial_x U_0(x)$; there exist $a, c > 0$ such that for all $x \in \mathbf{R}$ $|U_0(x)| \leq a(1 + |x|)$ and $|U'_0(x)| \leq c \exp(a|x|)$. These conditions are satisfied when u_0 is uniformly bounded. They include also some initial data with wild oscillations at infinity, e.g. $u_0(x) = e^x \sin e^x$ is allowed.

We first state the results concerning the linear equation (1.12). We note that the stochastic integral in that equation, according to (2.5), is well defined, in $L^2(dP^w)$, if

$$\begin{aligned} & \left\| \int_0^t G_{t-s} * (\psi_s dW_s)(x) \right\|_2^2 \\ &= \mathbf{E}^w \int_0^t ds \int dy dy' G_{t-s}(x-y) G_{t-s}(x-y') \\ & \quad \times C(y, y') \psi_s(y) \psi_s(y') < \infty. \end{aligned} \tag{2.11}$$

We now introduce an auxiliary brownian motion which will permit to write a solution of (1.12) as a generalized Feynman-Kac formula. Let $dP_{x,t}^\beta = dy dP_{y,0;x,t}^\beta$, where $dP_{y,0;x,t}^\beta$ is the measure of a backward brownian motion with diffusion coefficient 2ν starting at time t in x and arriving at time 0 in y . We will denote by $\mathbf{E}_{x,t}^\beta$ the expectation with respect to the probability measure $dP_{x,t}^\beta$. We stress that the brownian motion β is independent from the brownian sheet W .

Proposition 2.1. *Let $\psi_0 = \exp \left\{ \frac{1}{2\nu} U_0(x) \right\}$, the assumptions on U_0 imply that $\psi_0 \in C^1(\mathbf{R})$ and $\exists c_1, c_2, a > 0$:*

$$\forall x \in \mathbf{R} \quad c_1 e^{-a|x|} \leq \psi_0(x) \leq c_2 e^{a|x|}, \quad |\psi'_0(x)| \leq c_2 e^{a|x|}. \tag{2.12}$$

Set

$$\psi_t(x) = \mathbf{E}_{x,t}^\beta (\psi_0(\beta_0) e^{-\frac{\varepsilon}{2\nu} \int_0^t dW_s(\beta_s)}), \tag{2.13}$$

then $\forall (t, x) \in [0, T] \times \mathbf{R}$,

- (i) $\forall p \geq 1 \psi_t(x) \in L^p(dP^w)$ and it satisfies condition (2.11).
- (ii) dP^w -a.s. $\psi_t(x)$ is a solution of (1.12); $t \mapsto \psi_t(x)$ is dP^w -a.s. locally Hölder continuous with exponent $\alpha < 1/2$.
- (iii) $x \mapsto \psi_t(x)$ is dP^w -a.s. differentiable; its derivative is

$$\partial_x \psi_t(x) = G'_t * \psi_0(x) + \frac{\varepsilon}{2\nu} \int_0^t G'_{t-s} * (V \psi_s ds - \psi_s dW_s)(x) \tag{2.14}$$

$\forall p \geq 1 \partial_x \psi_t(x) \in L^p(dP^w)$. Furthermore the application $(t, x) \mapsto \partial_x \psi_t(x)$ is dP^w -a.s. locally Hölder continuous with exponent $\alpha < 1/2$ in space and $\alpha < 1/4$ in time.

(iv) dP^w - a.s. $\psi_t(x) > 0$.
 The properties (i)–(ii) are statements about the existence of the solution of (1.12), (iii)–(iv) are required to perform the Cole-Hopf transformation.

From Proposition 2.1 we have the following theorem which states the existence result for the stochastic Burgers equation (1.5).

Theorem 2.2. Set $u_t(x) = -2\nu \partial_x \log \psi_t(x)$, with $\psi_t(x)$ given by (2.13).

Then $u_t = u_t(x)$ is a $C^0(\mathbf{R})$ -valued, \mathcal{F}_t -adapted process, such that for all $(t, x) \in [0, T] \times \mathbf{R}$,

- (i) $\forall p \geq 1 u_t(x) \in L^p(dP^w)$; $(t, x) \mapsto u_t(x)$ is dP^w -a.s. locally Hölder continuous with exponent $\alpha < 1/2$ in space and $\alpha < 1/4$ in time.
- (ii) $u_t(x)$ solves (1.5) dP^w -a.s.

The paper is structured as follows. In the next section we prove a regularizing property for the stochastic convolution with the heat kernel; this is the main technical result in the paper. In Sect. 4 we prove, assuming Proposition 2.1, Theorem 2.2. Proposition 2.1 is then proved in Sect. 5. Moment estimates for $\psi_t(x)$ and $\log \psi_t(x)$ are obtained in Sect. 6; they give some insight on the behaviour of $u_t(x)$ at large x . Finally in Sect. 7 some open problems are discussed.

To simplify the notations we assume $\varepsilon = 1, \nu = 1/2$.

3. Stochastic Convolution with the Heat Kernel

We here extend to the stochastic case the regularization property of the heat kernel. In particular we show that the brownian sheet integral of the convolution of the heat kernel with a locally Hölder continuous process is dP^w -a.s. differentiable. The derivative is dP^w -a.s. locally Hölder continuous. This is the result that permits us to prove the differentiability of $x \mapsto \psi_t(x)$ in Proposition 2.1, hence the Cole-Hopf transformation is meaningful also for the stochastic Burgers equation. This result, however, has an independent interest.

Theorem 3.1. Let $\xi_t = \xi_t(x)$ a continuous \mathcal{F}_t -adapted process, assume the following properties hold for some $p \geq 2$:

- (a) $\exists c_1, a_1 > 0$ such that $\forall x \in \mathbf{R} \sup_{t \in [0, T]} \|\xi_t(x)\|_p \leq c_1 e^{a_1|x|}$.
- (b) $\exists c_2, a_2, \alpha > 0$ such that

$$\forall x, y \in \mathbf{R}, \forall t \in [0, T] \quad \|\xi_t(y) - \xi_t(x)\|_p \leq c_2 e^{a_2(|x|+|y|)} |x - y|^\alpha. \tag{3.1}$$

Define

$$F_t(x) = \int_0^t G_{t-s} * (\xi_s dW_s)(x). \tag{3.2}$$

Then

(i) $x \mapsto F_t(x)$ is $L^p(dP^w)$ differentiable; its derivative is

$$\partial_x F_t(x) = \int_0^t G'_{t-s} * (\xi_s dW_s)(x) \tag{3.3}$$

and $\exists c'_1, a'_1 > 0$ such that $\forall x \in \mathbf{R} \sup_{t \in [0, T]} \|\partial_x F_t(x)\|_p \leq c'_1 e^{a'_1 |x|}$.

(ii) $x \mapsto \partial_x F_t(x)$ is $L^p(dP^w)$ locally Hölder continuous with exponent less than α , i.e. for any $\varepsilon > 0$,

$$\forall x, y \in \mathbf{R}, \forall t \in [0, T] \quad \|\partial_y F_t(y) - \partial_x F_t(x)\|_p \leq c'_2 e^{\alpha_2'(|x|+|y|)} |x - y|^{\alpha - \varepsilon}. \tag{3.4}$$

(iii) If the conditions (a)–(b) are satisfied for all $p < \infty$, in general with c and a p -dependent constants, then $x \mapsto F_t(x)$ is also dP^w -a.s. differentiable; $x \mapsto \partial_x F_t(x)$ is dP^w -a.s. Hölder continuous with exponent smaller than α .

Proof. (i) We first discuss in detail the case $p = 2$. Let us introduce the notations $\Delta_h f(x) = h^{-1}(f(x + h) - f(x))$ and $R_f(x, h) = \Delta_h f(x) - \partial_x f(x)$; we will show that

$$\begin{aligned} & \left\| \int_0^t R_{G_{t-s}}(h) * (\xi_s dW_s)(x) \right\|_2^2 \\ &= \mathbf{E}^w \int_0^t ds \int dy dy' R_{G_{t-s}}(x - y, h) \\ & \quad \times R_{G_{t-s}}(x - y', h) C(y, y') \xi_s(y) \xi_s(y') \end{aligned} \tag{3.5}$$

converges to 0 when $h \rightarrow 0$. This requires an exchange of the limit $h \rightarrow 0$ with the integrals; due the specificity of this problem we cannot appeal to general theorems but we will estimate explicitly the integral.

Trying to bound directly the integrand in (3.5) a non-integrable singularity $(t - s)^{-1}$ appears. We thus need to exhibit a cancellation.

The key point is that $\int dy R_{G_{t-s}}(x - y, h) = 0$, so we can replace the right-hand side of (3.5) by

$$\int_0^t ds \int dy dy' R_{G_{t-s}}(x - y, h) R_{G_{t-s}}(x - y', h) \mathbf{E}^w(\Gamma_s(x, y, y')), \tag{3.6}$$

where

$$\Gamma_s(x, y, y') = C(y, y') \xi_s(y) \xi_s(y') - C(y', x) \xi_s(y') \xi_s(x). \tag{3.7}$$

We can now use the Hölder continuity assumption. In fact from the hypothesis (a), (b) and the global Lipschitz property for $C(y, y')$, we get the bound

$$\sup_{s \in [0, T]} \mathbf{E}^w |\Gamma_s(x, y, y')| \leq c e^{\alpha(|y|+|y'|+|x|)} |y - x|^\alpha. \tag{3.8}$$

Let us first consider in (3.6) the integral in dy' , we show it can be bounded by $(t-s)^{-1/2}$. We have

$$\int_{|y'-x|\leq 2|h|} dy' R_{G_{t-s}}(x-y', h) e^{a|y'|} \leq c e^{a(2|h|+|x|)} \frac{1}{\sqrt{t-s}}; \quad (3.9)$$

on the other hand, using the Lagrange theorem we get

$$\begin{aligned} & \int_{|y'-x|>2|h|} dy' \Delta_h G_{t-s}(x-y') e^{a|y'|} \\ & \leq c \int_{|z|>2|h|} dz \frac{|z+\zeta|}{(t-s)^{3/2}} e^{-\frac{(z+\zeta)^2}{2(t-s)}} e^{a(|z|+|x|)} \\ & \leq c e^{a|x|} \int_{|z|>2|h|} dz \frac{|z|+|h|}{(t-s)^{3/2}} e^{-\frac{z^2}{8(t-s)}} e^{a|z|} \leq c e^{a|z|} \frac{1}{\sqrt{t-s}}, \end{aligned} \quad (3.10)$$

where $|\zeta| \leq |h|$ and thus $|z+\zeta| > \frac{1}{2}|z|$ for $|z| > 2|h|$.

We can proceed analogously for the term with $\partial_x G_{t-s}(x-y)$.

Using the previous bound and the Hölder property, we now prove that (3.6) vanishes when $h \rightarrow 0$. Due to (3.9) and (3.10) it is enough to show that

$$\lim_{h \rightarrow 0} \int_0^t ds \frac{1}{\sqrt{t-s}} \int dy R_{G_{t-s}}(x-y, h) e^{a|y|} |y-x|^\alpha = 0. \quad (3.11)$$

We consider first the region $|x-y| \leq 2|h|$. To prove (3.11) in this region we show that the integrals arising from each of the three terms in R_G converge to zero. We have

$$\begin{aligned} & (t-s)^{-1/2} \int_{|y-x|\leq 2|h|} dy |\partial_x G_{t-s}(x-y)| e^{a|y|} |y-x|^\alpha \\ & \leq c e^{a|x|} \int_{|z|\leq 2|h|} dz \frac{|z|^{1+\alpha}}{(t-s)^{3/2}} G_{t-s}(z) \leq c e^{a|x|} h^{\alpha/2} (t-s)^{-1+\alpha/4} \end{aligned} \quad (3.12)$$

which is integrable in ds . Analogously

$$\begin{aligned} & (t-s)^{-1/2} \int_{|y-x|\leq 2|h|} dy |h^{-1} G_{t-s}(x-y)| e^{a|y|} |y-x|^\alpha \\ & \leq c e^{a|x|} |h|^{-1+\alpha/2} (t-s)^{-1+\alpha/2} \int_{|z|\leq 2|h|} dz \frac{|z|^{\alpha/2}}{(t-s)^{\alpha/4}} e^{-\frac{z^2}{2(t-s)}} \\ & \leq c e^{a|x|} |h|^{\alpha/2} (t-s)^{-1+\alpha/4}. \end{aligned} \quad (3.13)$$

For the last term we use the following estimate on the heat kernel:

$$G_t(z) \leq \frac{c}{\sqrt{t+|z|}}, \quad (3.14)$$

and integrate explicitly on ds . We have

$$\begin{aligned} & \int_0^t ds (t-s)^{-1/2} \int_{|y-x| \leq 2|h|} dy |h|^{-1} G_{t-s}(x+h-y) e^{a|y|} |y-x|^\alpha \\ & \leq ce^{a|x|} \int_{|z| \leq 2|h|} dz \int_0^t ds (t-s)^{-1/2} |h|^{-1} |z|^\alpha \frac{c}{\sqrt{t-s+|z+h|}} \\ & = ce^{a|x|} \int_{|z| \leq 2|h|} dz |h|^{-1} |z|^\alpha \log \left(1 + \frac{\sqrt{t}}{|z+h|} \right) \leq ce^{a|x|} c(b) |h|^{\alpha-b} \end{aligned} \quad (3.15)$$

for any $b > 0$ since the logarithm is bounded by any power. Then it is enough to take $b < \alpha$.

We now consider the case $|x-y| > 2|h|$; we show that the integrand is uniformly bounded; we are then allowed to take the limit $h \rightarrow 0$ inside the integrals and (3.11) follows. We have

$$\begin{aligned} & (t-s)^{-1/2} \int_{|y-x| > 2|h|} dy |\partial_x G_{t-s}(x-y)| e^{a|y|} |y-x|^\alpha \\ & \leq ce^{a|x|} \int dz \frac{|z|^{1+\alpha}}{(t-s)^{3/2}} G_{t-s}(z) e^{a|z|} \leq ce^{a|x|} (t-s)^{-1+\alpha/2} \end{aligned} \quad (3.16)$$

and, proceeding as in (3.10),

$$\begin{aligned} & \int_{|y'-x| > 2|h|} dy' \Delta_h G_{t-s}(x-y') e^{a|y'|} \\ & \leq ce^{a|x|} \int_{|z| > 2|h|} dz \frac{|z|^{1+\alpha}}{(t-s)^{3/2}} e^{-\frac{z^2}{8(t-s)}} e^{a|z|} \end{aligned} \quad (3.17)$$

(3.16) and (3.17) are integrable in ds .

This concludes the proof of the $L^2(dP^w)$ differentiability of $x \mapsto F_t(x)$.

The case $p > 2$ can be treated essentially in the same way. In fact from the B.D.G. inequality (2.6) we get

$$\begin{aligned} & \left\| \int_0^t R_{G_{t-s}}(h) * (\xi_s dW_s)(x) \right\|_p \leq c(p) \left\| \int_0^t ds \int dy dy' R_{G_{t-s}}(x-y, h) \right. \\ & \quad \left. \times R_{G_{t-s}}(x-y', h) C(y, y') \xi_s(y) \xi_s(y') \right\|_{p/2}^{1/2} \\ & \leq c(p) \left[\int_0^t ds \int dy dy' |R_{G_{t-s}}(x-y, h)| \right. \\ & \quad \left. \times |R_{G_{t-s}}(x-y', h)| \|\Gamma_s(x, y, y')\|_{p/2} \right]^{1/2}. \end{aligned} \quad (3.18)$$

The proof now continues as for $p = 2$. We remark that, due to the explicit dependence on t of the heat kernel, $F_t(x)$ defined in (3.2) is not a martingale; however the B.D.G. inequality holds as follows from the following argument. Consider first

$$\int_0^t G_{t_1-s} * (\xi_s dW_s)(x) \quad \text{for } t < t_1 \tag{3.19}$$

which is a \mathcal{F}_t martingale; apply the B.D.G. inequality and then let $t \rightarrow t_1$.

(ii) It follows exactly the same steps of (i), but with $\Delta_h G_{t-s}(x-y) - \partial_x G_{t-s}(x-y)$ in (3.5) replaced by $\partial_x G_{t-s}(x+h-y) - \partial_x G_{t-s}(x-y)$.

(iii) From the above discussion we have

$$\forall p \geq 1 \quad \exists c: \forall \varepsilon > 0 \quad \|\Delta_h \psi_t(x) - \partial_x \psi_t(x)\|_p \leq ch^{\alpha-\varepsilon} \tag{3.20}$$

the dP^w -a.s. differentiability follows then from Kolmogorov Theorem (see e.g. [13], I2.1). Analogously the dP^w -a.s. Hölder continuity of $x \mapsto \partial_x F_t(x)$ follows from (ii). \square

4. Cole-Hopf Transformation and Ito Calculus

In this section we show how Theorem 2.2 follows from Proposition 2.1. In order to verify that the process $u_t(x) = -\partial_x \log \psi_t(x)$ satisfies Burgers equation (1.5) we introduce a regularization. This procedure is needed to apply stochastic calculus. After the regularization is removed we obtain a mild form of the Ito formula for a function of the process $\psi_t(x)$. Finally we show how translation invariance for Burgers equation is recovered via the Cole-Hopf transformation.

Proof of Theorem 2.2. (i) From Jensen inequality and (2.13) we have

$$\frac{1}{\psi_t(x)} = \frac{1}{\mathbf{E}_{x,t}^\beta(\psi_0(\beta_0) e^{-\int_0^t dW_s(\beta_s)})} \leq \mathbf{E}_{x,t}^\beta(\psi_0(\beta_0)^{-1} e^{\int_0^t dW_s(\beta_s)}). \tag{4.1}$$

As it will be done in the proof of (i) in Proposition 2.1 it can be verified that the right-hand side of (4.1) is in $L^p(dP^w)$. From this and (iii) of Proposition 2.1 it follows that $u_t(x) \in L^p(dP^w)$. The Hölder continuity follows directly from statements (ii)–(iv) in Proposition 2.1.

(ii) Let us introduce a regularization of the brownian sheet

$$W_s^\kappa(x) = \delta^\kappa * W_s(x) = \int dy \delta^\kappa(x-y) W_s(y), \tag{4.2}$$

where $\delta^\kappa(x) = \kappa h(\kappa x)$ with $h \in C_0^\infty(\mathbf{R})$ a smooth positive function with compact support and $\int dx h(x) = 1$. The covariance of the process (4.2) is $t \wedge t' C^\kappa(x, x')$, where $C^\kappa = \delta^\kappa * C * \delta^\kappa$. Correspondingly we have $\psi_t^\kappa(x)$ which satisfies the regularized version of (1.11), i.e.

$$\begin{aligned} d\psi_t^\kappa(x) &= \left(\frac{1}{2} \partial_x^2 \psi_t^\kappa(x) + V^\kappa(x) \psi_t^\kappa(x)\right) dt - \psi_t^\kappa(x) dW_t^\kappa(x), \\ \psi_{t=0}^\kappa(x) &= \psi_0(x), \end{aligned} \tag{4.3}$$

where $V^\kappa(x) = \frac{1}{2} C^\kappa(x, x)$. We remark that the second derivative with respect to x is now meaningful.

Let us establish the convergence of the regularized process:

$$\lim_{\kappa \rightarrow \infty} \|\psi_t^\kappa(x) - \psi_t(x)\|_p = 0. \tag{4.4}$$

Using the Feynman-Kac representation (2.13) and denoting by $M_\beta^\kappa(t)$ the curvilinear stochastic integral for the regularized brownian sheet we have

$$\begin{aligned} & \|\psi_t^\kappa(x) - \psi_t(x)\|_p \\ & \leq \|\mathbf{E}_{x,t}^\beta(\psi_0(\beta_0)(e^{-M_\beta^\kappa(t)} + e^{-M_\beta(t)})|M_\beta^\kappa(t) - M_\beta(t)|)\|_p \\ & \leq \mathbf{E}_{x,t}^\beta(\psi_0(\beta_0)) \|e^{-M_\beta^\kappa(t)} + e^{-M_\beta(t)}\|_{2p} \|M_\beta^\kappa(t) - M_\beta(t)\|_{2p}. \end{aligned} \tag{4.5}$$

From the bound $|C * \delta^\kappa(x, y) - C(x, y)| \leq c\kappa^{-1}$ we see that the second factor converges, uniformly in β , to 0 as $\kappa \rightarrow \infty$. From Proposition 2.1, the first factor is bounded uniformly in κ .

Using the expression (2.14) for $\partial_x \psi_t(x)$ and following the same argument of Theorem 3.1, it is easy to prove also the $L^p(dP^w)$ convergence of the derivative, i.e.

$$\lim_{\kappa \rightarrow \infty} \|\partial_x \psi_t^\kappa(x) - \partial_x \psi_t(x)\|_p = 0. \tag{4.6}$$

Let $u_t^\kappa(x) = -\partial_x \log \psi_t^\kappa(x)$: we show it satisfies

$$u_t^\kappa(x) = G_t * u_0(x) - \frac{1}{2} \int_0^t ds G'_{t-s} * (u_s^\kappa)^2(x) + \int_0^t G'_{t-s} * dW_s^\kappa(x). \tag{4.7}$$

Integrating by parts we have the following identity:

$$\begin{aligned} & \int_0^t ds G_{t-s} * \left(\frac{\partial^2 \psi_s^\kappa}{\psi_s^\kappa} \right) (x) \\ & = \int_0^t ds G'_{t-s} * \left(\frac{\partial \psi_s^\kappa}{\psi_s^\kappa} \right) (x) + \int_0^t ds G_{t-s} * \left(\frac{\partial \psi_s^\kappa}{\psi_s^\kappa} \right)^2 (x), \end{aligned} \tag{4.8}$$

where ∂ means the derivative with respect to the integrated space variable.

Using (4.3) it is a simple Ito calculus exercise to compute the stochastic differential (with respect to s) of $G_{t-s}(x - y) \log \psi_s^\kappa(y)$. By (4.8) one then obtains

$$\begin{aligned} \log \psi_t^\kappa(x) &= G_t * \log \psi_0(x) + \frac{1}{2} \int_0^t ds G_{t-s} * \left(\frac{\partial \psi_s^\kappa}{\psi_s^\kappa} \right)^2 (x) \\ & \quad - \int_0^t G_{t-s} * dW_s^\kappa(x). \end{aligned} \tag{4.9}$$

Differentiating the above identity we verify that (4.7) is satisfied. We stress that the equation satisfied by $u_t^\kappa(x)$, defined through the Cole-Hopf transformation, would not have been (4.7) if we had interpreted (1.9) in the Ito sense.

We note that from (4.4) and (4.6) the $L^p(dP^w)$ convergence of the right-hand side of (4.8) follows, as $\kappa \rightarrow \infty$. That limit defines $\partial_x^2 \psi_s(x) / \psi_s(x)$ when convoluted with G_{t-s} . In fact, whereas $\partial_x \psi_t(x)$ exists, as stated in Proposition 2.1, the second

derivative of $\psi_t(x)$ does not have a meaning in the classical sense. Moreover each term in (4.9) converges to the non-regularized one. In the limit $\kappa \rightarrow \infty$ (4.9) is then a mild Ito formula.

In order to complete the proof of Theorem 2.2 we now take the limit $\kappa \rightarrow \infty$ in (4.7) and show that each term converges, in $L^2(dP^w)$, to the corresponding non-regularized one in (1.5).

Let us discuss first the stochastic integrals. Along the same line of Theorem 3.1 it can be easily proved that

$$\lim_{\kappa \rightarrow \infty} \left\| \int_0^t G'_{t-s} * (dW_s^\kappa - dW_s)(x) \right\|_2 = 0. \tag{4.10}$$

We now consider

$$\begin{aligned} \|u_t(x) - u_t^\kappa(x)\|_2 &\leq \left\| \frac{\partial_x \psi_t^\kappa(x)}{\psi_t(x)\psi_t^\kappa(x)} \right\|_4 \|\psi_t(x) - \psi_t^\kappa(x)\|_4 \\ &\quad + \left\| \frac{1}{\psi_t(x)} \right\|_4 \|\partial_x \psi_t(x) - \partial_x \psi_t^\kappa(x)\|_4, \end{aligned} \tag{4.11}$$

which converges to 0 as $\kappa \rightarrow \infty$ by (4.4), (4.6) and (i).

For the last term we have

$$\begin{aligned} &\left\| \int_0^t G'_{t-s} * ((u_s^\kappa)^2 - (u_s)^2)(x) \right\|_2 \\ &\leq \int_0^t ds \int dy |\partial_x G_{t-s}(x-y)| \|u_s^\kappa(y) + u_s(y)\|_4 \|u_s^\kappa(y) - u_s(y)\|_4, \end{aligned} \tag{4.12}$$

and $\|u_s^\kappa(y) - u_s(y)\|_4$ can be bounded as in (4.11) by a uniformly integrable function vanishing in the limit $\kappa \rightarrow \infty$.

This implies statement (ii) of the theorem: let us write Eq. (4.7) and (1.5) in the form $F^\kappa(u^\kappa) = 0$ and $F(u) = 0$. We have

$$\|F(u)\|_2 \leq \|F(u) - F^\kappa(u^\kappa)\|_2 + \|F^\kappa(u^\kappa)\|_2; \tag{4.13}$$

the first term on the right-hand side vanishes as $\kappa \rightarrow \infty$, the second one is identically 0. \square

Translation Invariance. We conclude this section showing how translation invariance for Burgers equation (1.5) is recovered. Let us define the brownian sheet with respect to a point $x = a$ instead of $x = 0$, i.e. $W_t^a(x) = W_t(x) - W_t(a)$, whose covariance is

$$\mathbf{E}^{w^a}(W_t^a(x)W_t^a(x')) = t \wedge t' \theta((x-a)(x'-a)) |x-a| \wedge |x'-a|. \tag{4.14}$$

The solution of (1.12) with $W_t(x)$ replaced by $W_t^a(x)$ is then

$$\psi_t^a(x) = \mathbf{E}_{x,t}^\beta(\psi_0(\beta_0) e^{-\int_0^t dW_s^a(\beta_s)}) = e^{-\int_0^t dW_s(a)} \mathbf{E}_{x,t}^\beta(\psi_0(\beta_0) e^{-\int_0^t dW_s(\beta_s)}). \tag{4.15}$$

In the above expression the dependence on a is factorized, so that, when the Cole-Hopf transformation is performed, it cancels out completely. We remark that the choice of a is related to the possibility to multiply $\psi_t(x)$ by an arbitrary time dependent factor which disappears when (4.9) is differentiated to obtain (4.7).

5. Solution of the Linear Equation

In this section we study the linear equation with multiplicative *half white* noise (1.11); we prove the existence of a solution for its mild form (1.12). This solution is constructed explicitly through a generalized Feynman-Kac formula (2.13). We then study the typical properties of the trajectories: they are strictly positive and differentiable with respect to x . The last property is a direct application of Theorem 3.1.

Proof of Proposition 2.1. (i) It is enough to assume $p \in \mathbf{N}$. Let $\vec{\beta} = \{\beta^i\}_{i=1}^n$ be a family of n independent brownian motions; then, under dP^w , $\{M_{\beta^i}(t)\}_{i=1}^n$ are mean zero Gaussian processes with covariance matrix given by

$$\mathbf{E}^w(M_{\beta^i}(t)M_{\beta^j}(t')) = \int_0^{t \wedge t'} ds C(\beta_s^i, \beta_s^j). \tag{5.1}$$

This is a direct consequence of the gaussian property of the system $\{M_{\beta^i}(t)\}_{i=1}^n$ and (2.10).

We have

$$\begin{aligned} \|\psi_t(x)\|_p^p &= \left\| \prod_{i=1}^p \mathbf{E}_{x,t}^{\beta^i}(\psi_0(\beta_0^i)e^{-M_{\beta^i}(t)}) \right\|_p^p = \mathbf{E}_{x,t}^{\vec{\beta}} \mathbf{E}^w \left(\prod_{i=1}^p \psi_0(\beta_0^i)e^{-M_{\beta^i}(t)} \right) \\ &= \mathbf{E}_{x,t}^{\vec{\beta}} \left(\prod_{i=1}^p \psi_0(\beta_0^i) \exp \left\{ \frac{1}{2} \sum_{i,j} \int_0^t ds C(\beta_s^i, \beta_s^j) \right\} \right) \\ &\leq c_2^p \left[\mathbf{E}_{0,t}^{\beta} \left(\exp \left\{ a|x + \beta_0| + \frac{p}{2} \int_0^t ds |x + \beta_s| \right\} \right) \right]^p \\ &\leq c_2^p \exp \left\{ \frac{1}{2} p^2 |x|t + ap|x| + \frac{1}{4!} ((pt + 2a)^3 - (2a)^3) \right\}. \end{aligned} \tag{5.2}$$

From the above bound it is easy to check that condition (2.11) is satisfied.

(ii) We now verify that $\psi_t(x)$ given by the Feynman-Kac formula (2.13) is the solution of (1.12). We have

$$\begin{aligned} &\int_0^t G_{t-s} * (V \psi_s ds - \psi_s dW_s)(x) \\ &= \int_0^t \int dy G_{t-s}(x-y) \mathbf{E}_{s,y}^{\beta}(\psi_0(\beta_0)e^{-M_{\beta}(s)})(V(y)ds - dW_s(y)), \end{aligned} \tag{5.3}$$

the kernel $G_{t-s}(x-y)$ can be interpreted as the transition probability of the brownian motion β_s . Hence, using the Markov property, (5.3) equals

$$\mathbf{E}_{x,t}^{\beta} \left(\psi_0(\beta_0) \int_0^t e^{-M_{\beta}(s)}(V(\beta_s) ds - dW_s(\beta_s)) \right). \tag{5.4}$$

Computing the stochastic differential (with respect to W) of $\exp\{-M_\beta(s)\}$ we get

$$\begin{aligned} d(e^{-M_\beta(s)}) &= e^{-M_\beta(s)} \left(-dM_\beta(s) + \frac{1}{2} d\langle M_\beta, M_\beta \rangle_s \right) \\ &= e^{-M_\beta(s)} (-dW_s(\beta_s) + V(\beta_s) ds). \end{aligned} \tag{5.5}$$

Thus (5.4) is equal to

$$\mathbf{E}_{x,t}^\beta(\psi_0(\beta_0) (e^{-\int_0^t dW_s(\beta_s)} - 1)) = \psi_t(x) - G_t * \psi_0(x), \tag{5.6}$$

and Eq. (1.12) is verified.

We now discuss the continuity of $(t, x) \mapsto \psi_t(x)$. Using the Feynman-Kac representation (2.13), we prove the following bound. For any $p \geq 1$, $\exists c, a > 0$ such that $\forall t \in [0, T]$

$$\|\psi_t(x) - \psi_t(y)\|_p \leq ce^{a(|x|+|y|)} |x - y|^{1/2}. \tag{5.7}$$

From the triangular and Cauchy-Schwartz inequalities we get

$$\begin{aligned} \|\psi_t(x) - \psi_t(y)\|_p &\leq \|\mathbf{E}_{0,t}^\beta((\psi_0(\beta_0 + x) - \psi_0(\beta_0 + y))e^{-M_{\beta+x}(t)})\|_p \\ &\quad + \|\mathbf{E}_{0,t}^\beta(\psi_0(\beta_0 + y) (e^{-M_{\beta+x}(t)} + e^{-M_{\beta+y}(t)}) |M_{\beta+x}(t) - M_{\beta+y}(t)|)\|_p \\ &\leq ce^{a(|x|+|y|)} |y - x| \\ &\quad + \mathbf{E}_{0,t}^\beta(\psi_0(\beta_0 + y) \|e^{-M_{\beta+x}(t)} + e^{-M_{\beta+y}(t)}\|_{2p} \\ &\quad \times \|M_{\beta+x}(t) - M_{\beta+y}(t)\|_{2p}); \end{aligned} \tag{5.8}$$

on the other hand, $\|M_{\beta+x}(t) - M_{\beta+y}(t)\|_{2p}$ can be bounded by

$$\begin{aligned} c \left[\int_0^t ds (C(\beta_s + x, \beta_s + x) - 2C(\beta_s + x, \beta_s + y) + C(\beta_s + y, \beta_s + y)) \right]^{1/2} \\ \leq ct^{1/2} |x - y|^{1/2}. \end{aligned} \tag{5.9}$$

Finally (5.7) follows from (5.2), (5.8) and (5.9).

From (1.12), we can obtain

$$\|\psi_t(x) - \psi_s(x)\|_p \leq ce^{a|x|} |t - s|^{1/2}. \tag{5.10}$$

We do not prove (5.10) because in the proof of (iii) an analogous, but technically harder, estimate for $\partial_x \psi_t(x)$ will be obtained.

Using the Kolmogorov Theorem, from (5.7) and (5.10), it follows the dP^w -a.s. local Hölder continuity of exponent $\alpha < 1/2$ for $(t, x) \mapsto \psi_t(x)$.

(iii) We now show, using (1.12), that $x \mapsto \psi_t(x)$ is $L^p(dP^w)$ and dP^w -a.s. differentiable; the derivative is given by (2.14).

Let $\Delta_h f(x) = h^{-1}(f(x+h) - f(x))$, then from (1.12) we get

$$\Delta_h \psi_t(x) = \Delta_h G_t * \psi_0(x) + \int_0^t \Delta_h G_{t-s} * (V\psi_s ds - \psi_s d_s W)(x) \tag{5.11}$$

and, for $\partial_x \psi_t(x)$ given by (2.14), we have

$$\begin{aligned} \|\Delta_h \psi_t(x) - \partial_x \psi_t(x)\|_p &\leq \|(\Delta_h G_t - G'_t) * \psi_0(x)\| \\ &\quad + \left\| \int_0^t ds (\Delta_h G_{t-s} - G'_{t-s}) * \psi_s(x) \right\|_p \\ &\quad + \left\| \int_0^t (\Delta_h G_{t-s} - G'_{t-s}) * (\psi_s dW_s)(x) \right\|_p. \end{aligned} \tag{5.12}$$

From the assumptions on the initial condition ψ_0 we have

$$\lim_{h \rightarrow 0} \Delta_h G_t * \psi_0(x) = G'_t * \psi_0(x). \tag{5.13}$$

In Theorem 3.1 we showed in detail the convergence to zero of the last term in (5.12); the second one is easier.

Let us discuss the continuity property of the derivative. From (3.4), repeating the above discussion, it is easy to see that $x \mapsto \partial_x \psi_t(x)$ is $L^p(dP^w)$ locally Hölder continuous with exponent $\alpha < 1/4$.

This implies for $x \mapsto \psi_t(x)$ a better estimate than (5.7): it is locally Lipschitz, i.e.

$$\|\psi_t(x) - \psi_t(y)\|_p \leq ce^{\alpha(|x|+|y|)}|x - y|. \tag{5.14}$$

Applying again Theorem 3.1 with this bound instead of (5.7) we prove that $x \mapsto \partial_x \psi_t(x)$ is $L^p(dP^w)$ locally Hölder continuous with exponent $\alpha < 1/2$,

$$\|\partial_x \psi_t(x) - \partial_y \psi_t(y)\|_p \leq ce^{\alpha(|x|+|y|)}|x - y|^{\frac{1}{2}-\epsilon}. \tag{5.15}$$

We now discuss the $L^p(dP^w)$ Hölder continuity of $t \mapsto \partial_x \psi_t(x)$. The bound (5.10) of (ii) is obtained in the same way. We consider $p = 2$, the general case follows via the B.D.G. inequality as in Theorem 3.1.

Let $k > 0$, from (2.14) we have, omitting the dependence on the space variable x ,

$$\begin{aligned} \|\partial_x \psi_{t+k} - \partial_x \psi_t\|_2 &\leq \|(G'_{t+k} - G'_t) * \psi_0| \\ &\quad + \left\| \int_0^{t+k} ds G'_{t+k-s} * (\psi_s V) - \int_0^t ds G'_{t-s} * (\psi_s V) \right\|_2 \\ &\quad + \left\| \int_0^{t+k} G'_{t+k-s} * (\psi_s dW_s) - \int_0^t G'_{t-s} * (\psi_s dW_s) \right\|_2. \end{aligned} \tag{5.16}$$

We consider in detail just the term with the stochastic integral in (5.16); we have

$$\begin{aligned} & \left\| \int_0^{t+k} G'_{t+k-s} * (\psi_s dW_s) - \int_0^t G'_{t-s} * (\psi_s dW_s) \right\|_2 \\ & \leq \left\| \int_t^{t+k} G'_{t+k-s} * (\psi_s dW_s) \right\|_2 + \left\| \int_{t-k}^t (G'_{t+k-s} - G'_{t-s}) * (\psi_s dW_s) \right\|_2 \\ & \quad + \left\| \int_0^{t-k} (G'_{t+k-s} - G'_{t-s}) * (\psi_s dW_s) \right\|_2, \end{aligned} \tag{5.17}$$

computing the $L^2(dP^w)$ norm, subtracting, as in (3.6), the appropriate null term

$$\mathbf{E}^w \int_t^{t+k} ds \int dy dy' \partial_x G_{t+k-s}(x-y) \partial_x G_{t+k-s}(x-y') C(y', x) \psi_s(y') \psi_s(x),$$

and using the Lipschitz property for $x \mapsto \psi_t(x)$, the first term in (5.17) can be bounded by

$$\begin{aligned} & ce^{a|x|} \left[\int_t^{t+k} ds \int dz dz' |\partial_z G_{t+k-s}(z)| |\partial_{z'} G_{t+k-s}(z')| |z| e^{a(|z|+|z'|)} \right]^{1/2} \\ & \leq ce^{a|x|} \left[\int_t^{t+k} ds (t+k-s)^{-1/2} \right]^{1/2} \leq ce^{a|x|} k^{1/4}. \end{aligned}$$

The second term in (5.17) is analogous, for the third term we have

$$\int dz' |\partial_{z'} G_{t+k-s}(z') - \partial_{z'} G_{t-s}(z')| e^{a|z'|} \leq c(t-s)^{-1/2},$$

and using Lagrange theorem

$$\begin{aligned} & \int_{z^2 > k} dz |\partial_z G_{t+k-s}(z) - \partial_z G_{t-s}(z)| |z| e^{a|z|} \\ & \leq c \int_{z^2 > k} dz k \frac{z^2}{(t-s+\sigma)^{5/2}} \left(3 + \frac{z^2}{t-s+\sigma} \right) e^{-\frac{z^2}{2(t-s+\sigma)}} e^{a|z|} \\ & \leq ck^{\frac{1}{2}-\varepsilon} \int dz |z|^{3+2\varepsilon} (t-s)^{-2} \left(3 + \frac{z^2}{t-s} \right) G_{2(t-s)}(z) e^{a|z|} \\ & \leq ck^{\frac{1}{2}-\varepsilon} (t-s)^{-\frac{1}{2}+\varepsilon}, \end{aligned}$$

as $\sigma \in [0, k]$ and $t - s \geq k$. On the other hand,

$$\begin{aligned} & \int_{z^2 \leq k} dz |\partial_z G_{t+k-s}(z) - \partial_z G_{t-s}(z)| |z| e^{a|z|} \\ & \leq ck^{\frac{1}{2}-\varepsilon} \int dz |z|^{1+2\varepsilon} \left(\frac{1}{t+k-s} G_{t+k-s}(z) + \frac{1}{t-s} G_{t-s}(z) \right) e^{a|z|} \\ & \leq ck^{\frac{1}{2}-\varepsilon} (t-s)^{-\frac{1}{2}+\varepsilon}, \end{aligned}$$

hence (5.17) is bounded by $c \exp(a|x|) k^{\frac{1}{4}-\varepsilon}$.

We have thus proved

$$\forall p \geq 1, \exists c > 0: \forall \varepsilon > 0$$

$$\|\partial_x \psi_{t+k}(x+h) - \partial_x \psi_t(x)\|_p \leq ce^{a|x|} (\sqrt{h^2+k^2})^{\frac{1}{4}-\varepsilon}. \tag{5.18}$$

From the Kolmogorov Theorem follows the dP^w -a.s. local Hölder continuity of $(t, x) \mapsto \partial_x \psi_t(x)$. The Hölder exponent is $\alpha < 1/4$ in time and $\alpha < 1/2$ in space.

(iv) We use the expression of $\psi_t(x)$ given by the Feynman-Kac formula (2.13). From the hypotheses on ψ_0 and Jensen inequality we have

$$\psi_t(x) \geq c_1 \exp \left\{ -\mathbf{E}_{x,t}^\beta \left(a|\beta_0| + \int_0^t dW_s(\beta_s) \right) \right\} \tag{5.19}$$

as

$$\mathbf{E}^w \left(\mathbf{E}_{x,t}^\beta \left(\int_0^t dW_s(\beta_s) \right) \right)^2 < \infty;$$

the exponent on the right-hand side of (5.19) is dP^w -a.s. finite, hence $\psi_t(x) > 0$ dP^w -a.s. \square .

6. Moment Estimates

In this section we conclude the analysis for the moments of $\psi_t(x)$ initiated in Sect. 5. We obtain a bound from below. We then estimate the moments of $\log \psi_t(x)$ which grow slower. They will give some insight on the behavior of $u_t(x)$ for large x .

Moment estimates for $\psi_t(x)$

Proposition 6.1. *Let $\psi_t(x)$ as in Proposition 2.1, $p \in \mathbf{N}$; then*

$$\begin{aligned} & c_1^p e^{\frac{1}{2} p^2 |x| t - ap|x| + \frac{1}{4t} ((pt-2a)^3 - (2a)^3)} e^{-cp^2 t^{3/2}} \\ & \leq \|\psi_t(x)\|_p^p \leq c_2^p e^{\frac{1}{2} p^2 |x| t + ap|x| + \frac{1}{4t} ((pt+2a)^3 - (2a)^3)} \end{aligned} \tag{6.1}$$

where a is the constant appearing in (2.12).

Proof. Formula (5.2) gives the estimate from above of the p^{th} moment of ψ . We now estimate from below $\|\psi\|_p^p$. We assume $\psi_0(x) \geq c > 0$, i.e. $U_0(x)$ uniformly bounded and we obtain (6.1) with $a = 0$. The general case is only more tedious.

From (5.2) we have

$$\begin{aligned} \|\psi(t, x)\|_p^p &= \mathbf{E}_{x,t}^{\vec{\beta}} \left(\psi_0(\beta_0^1) \dots \psi_0(\beta_0^p) \exp \left\{ \frac{1}{2} \int_0^t ds \sum_{i,j} C(\beta_s^i, \beta_s^j) \right\} \right) \\ &\geq c^p \mathbf{E}_{x,t}^{\vec{\beta}} \left(\exp \left\{ \frac{p}{2} \int_0^t ds \sum_{i=1}^p \beta_s^i - \int_0^t ds \sum_{1 \leq i < j \leq p} |\beta_s^i - \beta_s^j| \right\} \right). \end{aligned} \tag{6.2}$$

We introduce the brownian motion $\gamma_s = \frac{1}{\sqrt{p}} \sum_{i=1}^p \beta_s^i$. The first term in the exponent of (6.2) is the integral of γ_s , the second one is independent on it; the expectation thus factorizes and we get

$$\|\psi(t, x)\|_p^p \geq c_1^p e^{\frac{1}{2} p^2 |x|t + \frac{1}{4!} p^3 t^3} e^{-cp^2 t^{3/2}} \quad \square \tag{6.3}$$

These results do not imply intermittence in the sense of [11] because the field is not spatially ergodic.

Moment estimates for log $\psi_t(x)$

We now study the moments of $\log \psi_t(x)$. We start with the estimate from above.

Proposition 6.2. *Let $\psi_t(x)$ as in Proposition 2.1, then for either t or $|x|$ large enough (i) for $p = 2n + 1, n \in \mathbf{N}$,*

$$\mathbf{E}^w(\log^p \psi_t(x)) \leq \left[c_1 + \frac{1}{2} |x|t + a|x| + \frac{1}{4!} ((t + 2a)^3 - (2a)^3) \right]^p, \tag{6.4}$$

(ii) for $p = 2n, n \in \mathbf{N}$,

$$\mathbf{E}^w(\log^p \psi_t(x)) \leq c_2^p \left[p + |x| (t + a) \frac{1}{4!} + ((t + 2a)^3 - (2a)^3) + \sqrt{p}((t|x|)^{3/4}) \right]^p. \tag{6.5}$$

Proof. (i) Let $L_p \in C^0(\mathbf{R}^+)$ be the convex hull of the function $\xi \mapsto \log^p \xi$, by Jensen inequality

$$\begin{aligned} \mathbf{E}^w(\log^p \psi_t(x)) &\leq \mathbf{E}^w(L_p(\psi_t(x))) \leq L_p(\mathbf{E}^w(\log^p \psi_t(x))) \\ &\leq L_p\left(ce^{\frac{1}{2} |x|t + a|x| + \frac{1}{4!} ((t+2a)^3 - (2a)^3)}\right) \end{aligned} \tag{6.6}$$

as L_p is an increasing function. We note that $L_p(\xi) = \log^p \xi$ for $\xi \geq \xi_p = \exp\{p - 1\}$. Thus, when either $|x|$ or t are large enough, we get the result.

(ii) For p even function $\xi \mapsto \log^p \xi$ is monotone decreasing for $\xi \in (0, 1]$, monotone increasing for $\xi > 1$ and concave for $\xi > \xi_p$. We bound separately the three regions. We have

$$\begin{aligned} \mathbf{E}^w(\chi_{\{\psi \leq 1\}} \log^p \psi_t(x)) &\leq \mathbf{E}^w \left(\mathbf{E}_{x,t}^{\beta} \left(\log \psi_0(\beta_0) - \int_0^t dW_s(\beta_s) \right) \right)^p \\ &\leq \left[\mathbf{E}_{x,t}^{\beta}(\log \psi_0(\beta_0)) + \left\| \mathbf{E}_{x,t}^{\beta} \left(\int_0^t dW_s(\beta_s) \right) \right\|_p \right]^p \\ &\leq [a(1 + |x|) + c\sqrt{p}((t|x|)^{1/2} + t^{3/4})]^p, \end{aligned} \tag{6.7}$$

where the last inequality is obtained by explicit computations of gaussian integrals.

On the other hand

$$\mathbf{E}^w(\chi_{\{1 < \psi \leq \xi_p\}} \log^p \psi_t(x)) \leq \log^p \xi_p. \tag{6.8}$$

Denote by $\tilde{\mathbf{E}}^w$ the expectation with respect to dP^w conditioned to the event $\{\psi > \xi_p\}$; by Jensen inequality we finally have

$$\begin{aligned} \mathbf{E}^w(\chi_{\{\psi > \xi_p\}} \log^p \psi_t(x)) &\leq P^w(\{\psi > \xi_p\}) \log \tilde{\mathbf{E}}^w(\psi_t(x)) \\ &\leq P^w(\{\psi > \xi_p\}) \log^p \frac{\mathbf{E}^w(\psi_t(x))}{P^w(\{\psi > \xi_p\})} \\ &\leq P^w(\psi > \xi_p) \left[\log c_2 + \frac{1}{2} |x| (t + a) \right. \\ &\quad \left. + \frac{1}{4!} ((t + 2a)^3 - (2a)^3) - \log P^w(\{\psi > \xi_p\}) \right]^p, \end{aligned} \tag{6.9}$$

where we used the bound in (6.1) for $p = 1$. Since for $\xi \in [0, 1]$, $\xi \log^p \xi \leq (c/e)^p$, (6.5) follows from (6.7), (6.8) and (6.9). \square

We now bound from below the moments of $\log \psi_t(x)$ under the following restrictions: there exist $c_1, c_2 > 0$ such that for all $x \in \mathbf{R}$, $1 < c_1 \leq \psi_0(x) \leq c_2$.

Proposition 6.3. *Let $\psi_t(x)$ as in Proposition 2.1, then*

(i) for $p = 2n + 1, n \in \mathbf{N}$,

$$\begin{aligned} \mathbf{E}^w(\log^p \psi_t(x)) &\geq \sqrt{p}! c^p [|x|(t \wedge (2x^2)) - (t \wedge (2x^2))^{3/2} \\ &\quad + \theta(t - 2x^2) \cdot (t^{3/2} - |x|^3)]^{\frac{p-1}{2}}. \end{aligned} \tag{6.10}$$

(ii) For $p = 2n, n \in \mathbf{N}$

$$\mathbf{E}^w(\log^p \psi_t(x)) \geq c_1^p ((p!)^3 (t^2 x^2 + t^3)^p)^{1/4} - c_2^p p! \tag{6.11}$$

Proof. (i) Since $\log^p \xi$ is increasing for p odd, by Jensen inequality we have

$$\begin{aligned} \mathbf{E}^w \log^p \psi_t(x) &\geq \mathbf{E}^w \left(\mathbf{E}_{x,t}^\beta \left(\log \psi_0(\beta_0) - \int_0^t dW_s(\beta_s) \right) \right)^p \\ &= \sum_{k=0}^p \frac{p!}{k!(p-k)!} (\mathbf{E}_{x,t}^\beta \log \psi_0(\beta_0))^{p-k} \cdot \mathbf{E}^w \left(\mathbf{E}_{x,t}^\beta \left(- \int_0^t dW_s(\beta_s) \right) \right)^k \\ &\geq p!! c^p \left[\mathbf{E}_{x,t}^{\beta^1, \beta^2} \int_0^t ds C(\beta_s^1, \beta_s^2) \right]^{\frac{p-1}{2}}. \end{aligned} \tag{6.12}$$

The bound (6.10) can then be obtained after an explicit computation of the expectation in (6.12).

(ii) Let $X = \psi_0(\beta_0) \exp\{-M_\beta(t)\}$, following the same argument of (6.9), we have

$$\begin{aligned} \mathbf{E}_{x,t}^\beta(\chi_{\{X > \xi_p\}} \log^p X) &\leq 2^{p-1} (\log^p \mathbf{E}_{x,t}^\beta(X) + (p/e)^p) \\ &= 2^{p-1} (\log^p \psi_t(x) + (p/e)^p) \end{aligned} \tag{6.13}$$

and

$$\mathbf{E}_{x,t}^\beta(\chi_{\{1 \leq X \leq \xi_p\}} \log^p X) \leq \log^p \xi_p = p - 1. \tag{6.14}$$

Thus

$$\begin{aligned} \log^p \psi_t(x) &\geq c_1^p \mathbf{E}_{x,t}^\beta \left(\chi_{\{X > 1\}} \sum_{k=0}^p \frac{p!}{k!(p-k)!} (-M_\beta(t))^k \log^{p-k} \psi_0(\beta_0) \right) - c_2^p p! \\ &\geq c_1^p \mathbf{E}_{x,t}^\beta \left(\chi_{\{X > 1\}} \sum_{k \text{ even}} \frac{p!}{k!(p-k)!} (-M_\beta(t))^k \log^{p-k} \psi_0(\beta_0) \right) \\ &\quad - c_2^p (p! + \mathbf{E}_{x,t}^\beta \log^p \psi_0(\beta_0)). \end{aligned} \tag{6.15}$$

As $\{X > 1\} \supset \{M_\beta(t) < 0\}$, taking the expectation with respect to dP^w of (6.15) the result follows. \square

As far as $u_t(x) = -\partial_x \psi_t(x) / \psi_t(x)$ is concerned, we have the following

Proposition 6.4. *Let $u_t(x)$ as in Theorem 2.2, then for almost all $t \in [0, T]$ and $\forall \delta > 0$,*

$$\liminf_{x \rightarrow \infty} \frac{\|u_t(x)\|_2}{|x|^{\frac{1}{2} + \delta}} = 0. \tag{6.16}$$

Proof. In Sect. 4 we obtained the identity

$$\log \psi_t(x) = G_t * \log \psi_0(x) + \frac{1}{2} \int_0^t ds G_{t-s} * u_s^2(x) - \int_0^t G_{t-s} dW_s(x), \tag{6.17}$$

taking the expectation with respect to dP^w , Proposition 6.2 implies

$$\int_0^t ds G_{t-s} * \|u_s\|_2^2(x) \leq c(1 + (t + a)|x| + (t + 2a)^3), \tag{6.18}$$

from which (6.16) is derived. \square

7. Concluding Remarks

We conclude with some remarks on open problems.

In this paper we have not discussed at all the questions of uniqueness of the solution we have constructed. This is not a simple problem. The difficulty comes from the unboundedness of the space domain. In the unperturbed case (1.1) uniqueness was obtained by Hopf [7], by reducing the problem to the uniqueness of positive solutions of the heat equation. In our case this approach presents various difficulties. In Sect. 4 we proved that to a class of solutions of the linear equation (1.12) can be associated a class of solutions of the Burgers equation (1.5). The converse statement is not obvious as the application of Ito calculus requires the introduction of a regularization which in this direction is more difficult to remove. Furthermore the uniqueness problem for (1.12) is difficult due to the bad behavior of the random potential at infinity. On the other hand, in an unbounded domain, a naive attempt to obtain a Gronwall inequality

for an appropriate norm of the solution of Eq. (1.5) fails due to the structure of the nonlinearity.

In the study of the stochastic Burgers equation a most relevant question is the behavior of the solutions when ν or ε or both tend to zero. Here we have to distinguish different regimes. We consider the solution over a given time interval.

- ν finite, $\varepsilon \rightarrow 0$. The study of this case amounts essentially to the construction of a large deviation theory for the Burger equation. The main difficulties in the construction of such a theory are again connected to the infinite domain.
- ε finite, $\nu \rightarrow 0$. A way to approach this problem at the formal level is to evaluate the Feynman-Kac formula by a Laplace type approximation. One obtains a formal expression which is the same one would obtain by solving the stochastically perturbed inviscid Burgers equation via the method of characteristics. This formal expression is a distribution valued process. This means that its square is not well defined and the mathematical interpretation of the stochastic inviscid equation is not apparent; some renormalization may be necessary. The relationship of these limit solutions, if they exist, to shock waves should be investigated. This problem has some similarity to that considered by Lax in [9].
- When $\varepsilon, \nu \rightarrow 0$, on the basis of heuristic arguments, the solution should converge to those of the unperturbed inviscid equation. However this conclusion may be false as the result may depend on the way the double limit is taken.

Acknowledgements. We thank R. S en or for useful discussions and the Ecole Polytechnique for hospitality at an early stage of this work, S. Olla for calling our attention to the Lax phenomenon and E. Presutti for helpful comments. This work was partially supported by EEC contract SC1*0394-C(EDB).

Note added in proof. After completing this work we received the following preprints, *The Burgers Equation with a Noisy Force* by H. Holden, T. Lindstr om, B.  ksendal, J. Ub oe and T.-S. Zhang, where the stochastic Burgers equation is studied in the framework of white noise calculus; *Stochastic Burgers Equation* by G. Da Prato, A. Debussche, R. Temam where a detailed study on a finite space interval is made with a different approach.

References

1. Albeverio, S., Molchanov, S.A., Surgailis, D.: Stratified Structure of the Universe and the Burgers' Equation – A Probabilistic Approach. Preprint (1993)
2. Benzi, R., Jona-Lasinio, G., Suter, A.: Stochastically Perturbed Landau-Ginzburg Equations. *J. Stat. Phys.* **55**, 505–522 (1989)
3. Burgers, J.M.: *The Nonlinear Diffusion Equation*. Dordrecht: D. Reidel, 1974
4. Butt , P.: Two dimensional Yang-Mills measure as a distribution valued brownian motion. SISSA Preprint 21/92/FM (1992)
5. Cole, J.D.: On a Quasi-Linear Parabolic Equation Occurring in Aerodynamics. *Quart. Appl. Math.* **9**, 225–236 (1951)
6. Gihkman, I.I., Skorohod, A.V.: *Stochastic Differential Equations*. Berlin Heidelberg New York: Springer 1972
7. Hopf, E.: The Partial Differential Equation $u_t + uu_x = \mu u_{xx}$. *Commun. Pure Appl. Math.* **3**, 201–230 (1950)
8. Ikeda, N., Watanabe, S.: *Stochastic Differential Equations and Diffusion Processes*. Amsterdam, Oxford, New York: North Holland Publishing, 1981
9. Lax, P.D.: The Zero Dispersion Limit, A Deterministic Analogue of Turbulence. *Commun. Pure Appl. Math.* **44**, 1047–1056 (1991)

10. Lighthill, M.J.: Viscosity Effects in Sound Waves of Finite Amplitude. In: Batchelor, G.K., Davies, R.M. (eds.) *Surveys in Mechanics*, Cambridge: Cambridge University Press, 1956, pp. 250–351
11. Molchanov, S.A.: Ideas in Theory of Random Media. *Acta Appl. Math.* **22**, 139–282 (1991) and references therein
12. Mueller, C.: On the Support of Solutions to the Heat Equation with Noise. *Stochastics and Stochastics Reports* **37**, 225–245 (1991)
13. Revuz, D., Yor, M.: *Continuous Martingales and Brownian Motion*. Berlin, Heidelberg, New York: Springer 1991
14. She, Z.S., Aurell, E., Frisch, U.: The Inviscid Burgers Equation with Initial Data of Brownian Type. *Commun. Math. Phys.* **148**, 623–641 (1992)
15. Sinai, Ya.G.: Two Results Concerning Asymptotic Behavior of Solution of the Burgers Equation with Force. *J. Stat. Phys.* **64**, 1–12 (1991)
16. Sinai, Ya.G.: Statistics of Shocks in Solutions of Inviscid Burgers Equation. *Commun. Math. Phys.* **148**, 601–621 (1992)
17. Walsh, J.B.: An introduction to stochastic partial differential equations. In: *Lecture Notes in Mathematics* **1180**. Berlin, Heidelberg, New York: Springer 1984

Communicated by Ya. G. Sinai